


SANIBEL

GENERAL

- SANIBEL is a very early variety (75 – 90 growing days)
- Sanibel makes a second skin, so the skin is every times red
- Take the seed out of the bag directly after arrival for better seed conditions
- Soil preparation; aim should be a loose topsoil
- Planting distance:

- Planting depth: normal


Aim of size	Seed size	Row distance	
		75 cm	90 cm
40/60	28/35	15-17 cm	13-14 cm
	35/55	27-30 cm	23-25 cm
60+	28/35	18-20 cm	15-17 cm
	35/55	32-34 cm	25-28 cm



- P_2O_5 80 kg/ha
- MgO 60 kg/ha
- Seed dressing: treatment is recommended; preferably furrow dressing (Azoxystrobin as a soil treatment shows good results against many fungi)

ATTENTION POINTS

- SANIBEL is sensitive for pre and post emergence application of Metribuzin (SENCOR), this should be avoided
- N-demand demand of SANIBEL is low, 20% less in comparison to other very early varieties
- N 140-180 kg/ha incl. N_{min} dependent from soil type/ location/packing size) – Splitting in two applications 3/4:1/4
- K_2O 300 - 350 kg/ha (as K_2SO_4) drought tolerance – with irrigation split of the amounts 1/2:1/2
- Sanibel is susceptible to leaf blight – early start and regularly treatments are recommended
- 1 or 2 treatments against aphids is recommended



All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.